Math 61 www.timetodare.com

1.2 Points, Lines, and Planes 1.3 Segments, Rays, and Angles

A Mathematical System consists of

1. Undefined Terms
- line
- plane
2. Defined Terms
3. Axioms or Postulates
4. Theorems

Set - a collection of objects.

A point

- the simplest geometric figure
- it is suggested by a dot on a piece of paper or by the tip of a pencil
- it has no size
- it has a position
- notation: uppercase italic letters A, B, C, etc.

A plane

- it is suggested by a flat surface with the idea that it extends without end in all directions
- notation: script letters P or Greek letters a, b, g

A line

- it is suggested by the edge of a table or a rail of a railroad track
- we think of it as determining the shortest distance between two points, as having no bends, as extending indefinitely in both directions
- notation: lowercase italic letters d, l, f, etc or using two points on it, as in \overrightarrow{AB}

After some simple terms such as "point", "line", and "plane" have been accepted as undefined, we can begin to define other terms by using them.

When is a statement a definition?

1. It names the term being defined.

2. It places the term into a set or category.

A good definition will possess these qualities: \{3. It distinguishes the defined term from other terms without providing unnecessary facts.

4. It is reversible.

Definition The set of all points is called **space**.

Definition Any set of points, lines, or planes in space is called a geometric figure.

Postulates

Geometry, or any deductive system, is very much like a game. Before playing the game, it is necessary to accept some basic rules, which we will call *postulates*. The postulates in geometry are man-made, just as the rules of football are, and what the subject will be like depends upon the nature of the postulates used. We will study the geometry called Euclidean, named after Euclid. For many centuries, it was the only geometry known, because it took man a long time to realize that more than one set of rules were possible.

Geometry has very few rules. We will need to supplement them with some of the rules of algebra with which you are already familiar. The rules, or postulates, of algebra concern numbers and operations performed on them.

<u>Properties of Equality</u> (1.2 – Postulates 1.6 – 1.12)

Reflexive Property Any real number is equal to itself.

a = a

Symmetric Property If a = b, then b = a

Transitive Property If a = b and b = c, then a = c

Addition Property If a = b, then a + c = b + c

a-c=b-c.

Multiplication Property If a = b, then $a \cdot c = b \cdot c$

 $\frac{a}{c} = \frac{b}{c}, \forall c \neq 0$.

Distributive Property a(b+c) = ab + ac

Substitution Property If a = b, then a can be substituted for b in any expression containing b.

The postulates of geometry deal with sets of points and their relationships.

Question Consider a single point. How many lines can pass through, or contain, it?

Question Now consider two points. How many lines can contain them?

Postulate 1:

Through two distinct points, there is exactly one line. (Two points determine a line.)

Definition

Points that lie on the same line are called collinear points.

F

Exercise #1

- a) Name three points that appear to be collinear.
- A [•]
- b) Name three points that appear to be noncollinear.

c ·

- c) How many lines can be drawn through point *A*?
- d) How many lines can be drawn through points A and B?
- e) How many lines can be drawn through points A, B, and C?

Exercise #2

- a) Make a drawing to illustrate three noncollinear points A, B, and C, and all of the lines they determine. How many lines are there in all?
- b) Make a drawing to illustrate four points, no three of which are collinear, and all of the lines they determine. How many lines are there in all?
- c) Make a drawing to illustrate five points, no three of which are collinear, and all of the lines they determine. How many lines are there in all?



d) Without making a drawing, can you figure out how many lines are determined by ten points, no three of which are collinear?

Postulate 2	2
-------------	---

Through three noncollinear points, there is exactly one plane. (Three noncollinear points determine a plane).

Definition

Points that lie in the same plane are called **coplanar points.**

Postulate 3

Given two distinct points in a plane, the line containing these points also lies in the plane.

Postulate 4

No plane contains all points in space.

Postulate 5

There is a one-to-one correspondence between the set of all points on a line and the set of all real numbers.

Exercise #3 (1.2 - #1,3)

Answer each question and make a drawing to illustrate each situation.

- 1. How many lines can be drawn between two distinct points?
- 2. If distinct points A and B are in a plane P, and point C is on the line determined by A and B, what can be said about point C?

Exercise #4 (1.2 - # 5, 6)

Identity the hypothesis and conclusion of each statement. Name the postulate illustrated by each statement.

Make a drawing to illustrate each situation.

A, B, and C are distinct points; l is a line; \boldsymbol{a} is a plane.

1. If A and B are on l, and A and B are on m, then m = l.

2. If A and B are on l, l is in \boldsymbol{a} , and C is on l, then C is in \boldsymbol{a} .

Exercise #5 (1.2 - #23)

Give the reasons that support each indicated step in the solution of the equation 3x + 2 = 4 + 5x

(1.2	25)	
State	ments	

Reasons

1.
$$3x + 2 = 4 + 5x$$

2.
$$3x+2-4=4-4+5x$$

3.
$$3x - 2 = 5x$$

4.
$$3x-3x-2=5x-3x$$

5.
$$-2 = 2x$$

6.
$$\frac{1}{2}(-2) = \frac{1}{2}(2x)$$

7.
$$-1 = x$$

8.
$$x = -1$$

Definition

A **line segment** is the part of a line that consists of two points (endpoints) and all points between them.

Question

Is the above definition a good definition?

Exercise #6

a) You have learned that the following statement is true:

If a statement is a definition, then its converse is true.

Does it necessarily follow that if its converse is not true, a statement cannot be a definition? Explain.

Decide which of the following true statements are good definitions of the italicized words by determining whether their converses are true.

- b) If something is *cold*, then it has a low temperature.
- c) A mandolin is a stringed musical instrument.
- d) A kitten is a young cat.
- e) An isosceles triangle is a triangle that has two congruent sides.

Note:

When both a statement and its converse are true, there is a convenient way to combine the two into one. It is by means of the phrase "if and only if".

Definition

The **distance** between two points is the length of the line segment \overline{AB} that joins the two points.

Example:

Draw two points and find the distance between them.

Postulate 6:

<u>Segment – Addition Postulate</u>

If B is a point of \overline{AC} and A - B - C, then AB + BC = AC

Exercise #7 | Given: M a point on \overline{AB}

AM = 2x+1 AB = 3x-2 AB = 4Find: x, AM, and MB.

Definition

Ray AB, denoted by \overrightarrow{AB} , consists of the point A (endpoint) together with all the points on the line \overrightarrow{AB} on the same side of A as B.


Definition

Two rays are **opposite rays** if they have a common endpoint and if their union is a straight line.

Exercise #8

In the figure, name:

- a) two opposite rays.
- b) two rays that are not opposite.



Definition

An **angle** is the union of two rays that share a common endpoint.

Example

Draw an angle, name it, and measure it.

Types of Angles

ACUTE ANGLE – an angle whose measure is less than 90° .

RIGHT ANGLE – an angle whose measure is exactly 90°.

OBTUSE ANGLE – an angle whose measure is between 90° and 180° .

STRAIGHT ANGLE – an angle whose sides form a line and whose measure is exactly 180°.


Exercise #9

Given the figure, which points lie

in the **interior** of $\angle BAC$?

on $\angle BAC$?

in the exterior of $\angle BAC$?



Postulate 7

Angle – Addition Postulate

If a point *P* lies in the interior of an angle ABC, then $m\angle ABP + m\angle PBC = m\angle ABC$

Classifying Pairs of Angles

Two angles are **adjacent angles** if they have a common vertex , share a common side, and have no interior points in common.

Two angles are **complementary** if their sum is 90° .

Two angle are supplementary if their sum is 180° .

When two lines intersect, the pairs of nonadjacent angles formed are known as vertical angles.

Example Draw two intersecting lines.

a) Which angles are vertical angles?

b) Which angles are supplementary?

Exercise #10 conclude


 $\angle FAC$ and $\angle CAD$ are adjacent and \overrightarrow{AF} and \overrightarrow{AD} are opposite rays. What can you

about $\angle FAC$ and $\angle CAD$?

Exercise #11

Given: $m\angle RST = 2x + 9$ $m\angle TSV = 3x - 2$ $m\angle RSV = 67^{\circ}$

Find: x.



Exercise #12 If $m \angle A = 27^{\circ}$ and $\angle A$ and $\angle B$ are complementary, find the measure of $\angle B$.